

Lnk International

Exhibits

012 3058

8074130



TRI-BUFFERED

Compare to the active ingredients in Bufferin®

Aspirin

Analgesic



PAIN RELIEVER • FEVER REDUCER

130 COATED BUFFERED ASPIRIN TABLETS
325 mg EACH



TRI-BUFFERED

Compare to the active ingredients in Bufferin®

Aspirin

Analgesic



PAIN RELIEVER • FEVER REDUCER

130 COATED BUFFERED ASPIRIN TABLETS
325 mg EACH



TRI-BUFFERED

Aspirin

Analgesic

PAIN RELIEVER • FEVER REDUCER

130 COATED BUFFERED ASPIRIN TABLETS
325 mg EACH



TRI-BUFFERED

Compare to the active ingredients in Bufferin®

Aspirin

Analgesic



PAIN RELIEVER • FEVER REDUCER

130 COATED BUFFERED ASPIRIN TABLETS
325 mg EACH

INDICATIONS: For relief of mild to moderate pain, headache, fever, and rheumatism. It is also used to reduce the risk of heart disease and stroke in certain patients.

CONTRAINDICATIONS: Do not take if you are allergic to aspirin or any of the ingredients. Do not take if you are taking other aspirin-containing products. Do not take if you have a history of stomach ulcers, bleeding disorders, or kidney disease. Do not take if you are pregnant or breastfeeding.

Warnings: Do not take with alcohol or other medications that may increase the risk of bleeding. Do not take if you have a history of asthma or other respiratory conditions. Do not take if you are taking blood thinners.

Directions: Take one tablet every 4 to 6 hours as needed. Do not take more than 6 tablets in 24 hours. Do not take for more than 10 days without consulting your doctor.

Side Effects: Common side effects include heartburn, stomach pain, and dizziness. Serious side effects include bleeding, stomach ulcers, and kidney damage.

Other Information: Keep out of reach of children. Store at room temperature. Do not use if the seal is broken or the tablets are discolored.

NDC 21849-100-1

ECKERD

Aspirin Plus

ASPIRIN, ALUMINA AND MAGNESIA TABLETS
ANALGESIC

PAIN RELIEF -- WITHOUT STOMACH DISCOMFORT

100 TABLETS

ECKERD

Aspirin Plus

ASPIRIN, ALUMINA AND MAGNESIA TABLETS
ANALGESIC

PAIN RELIEF -- WITHOUT STOMACH DISCOMFORT

100 TABLETS

TAMPER EVIDENCE: DO NOT USE IF IMPROPER SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

ECKERD

Aspirin Plus

DATE



ECKERD ITEM 508570
NDC, 19466-500-1

DRUG FACTS

Active Ingredients (In each tablet) Aspirin 5 grains, Magnesium Oxide 1/4 grain, Aluminum Oxide 1/4 grain.

Warnings
This product should not be taken by individuals with a known hypersensitivity to aspirin, salicylates, or other non-steroidal anti-inflammatory drugs (NSAIDs). It should not be taken by individuals with a history of asthma, urticaria, or allergic reactions to other NSAIDs. It should not be taken by individuals with a history of peptic ulcer disease, bleeding disorders, or kidney disease. It should not be taken by individuals who are taking other NSAIDs, anticoagulants, or antiplatelet drugs. It should not be taken by individuals who are taking other drugs that may increase the risk of bleeding. It should not be taken by individuals who are taking other drugs that may increase the risk of kidney disease. It should not be taken by individuals who are taking other drugs that may increase the risk of liver disease. It should not be taken by individuals who are taking other drugs that may increase the risk of heart disease. It should not be taken by individuals who are taking other drugs that may increase the risk of stroke. It should not be taken by individuals who are taking other drugs that may increase the risk of cancer. It should not be taken by individuals who are taking other drugs that may increase the risk of infection. It should not be taken by individuals who are taking other drugs that may increase the risk of other serious conditions.

Directions
Take 1 or 2 tablets 3 or 4 times a day as directed. Do not exceed the recommended dosage. Do not take this product if you are taking other NSAIDs, anticoagulants, or antiplatelet drugs. Do not take this product if you are taking other drugs that may increase the risk of bleeding, kidney disease, liver disease, heart disease, or stroke. Do not take this product if you are taking other drugs that may increase the risk of cancer, infection, or other serious conditions.

Other Information
This product contains aspirin, which may increase the risk of bleeding, kidney disease, liver disease, heart disease, or stroke. It may also increase the risk of cancer, infection, or other serious conditions. It should be used with caution in individuals who are taking other NSAIDs, anticoagulants, or antiplatelet drugs. It should be used with caution in individuals who are taking other drugs that may increase the risk of bleeding, kidney disease, liver disease, heart disease, or stroke. It should be used with caution in individuals who are taking other drugs that may increase the risk of cancer, infection, or other serious conditions.

DRUG FACTS

Warnings
This product should not be taken by individuals with a known hypersensitivity to aspirin, salicylates, or other non-steroidal anti-inflammatory drugs (NSAIDs). It should not be taken by individuals with a history of asthma, urticaria, or allergic reactions to other NSAIDs. It should not be taken by individuals with a history of peptic ulcer disease, bleeding disorders, or kidney disease. It should not be taken by individuals who are taking other NSAIDs, anticoagulants, or antiplatelet drugs. It should not be taken by individuals who are taking other drugs that may increase the risk of bleeding. It should not be taken by individuals who are taking other drugs that may increase the risk of kidney disease. It should not be taken by individuals who are taking other drugs that may increase the risk of liver disease. It should not be taken by individuals who are taking other drugs that may increase the risk of heart disease. It should not be taken by individuals who are taking other drugs that may increase the risk of stroke. It should not be taken by individuals who are taking other drugs that may increase the risk of cancer. It should not be taken by individuals who are taking other drugs that may increase the risk of infection. It should not be taken by individuals who are taking other drugs that may increase the risk of other serious conditions.

Directions
Take 1 or 2 tablets 3 or 4 times a day as directed. Do not exceed the recommended dosage. Do not take this product if you are taking other NSAIDs, anticoagulants, or antiplatelet drugs. Do not take this product if you are taking other drugs that may increase the risk of bleeding, kidney disease, liver disease, heart disease, or stroke. Do not take this product if you are taking other drugs that may increase the risk of cancer, infection, or other serious conditions.

Other Information
This product contains aspirin, which may increase the risk of bleeding, kidney disease, liver disease, heart disease, or stroke. It may also increase the risk of cancer, infection, or other serious conditions. It should be used with caution in individuals who are taking other NSAIDs, anticoagulants, or antiplatelet drugs. It should be used with caution in individuals who are taking other drugs that may increase the risk of bleeding, kidney disease, liver disease, heart disease, or stroke. It should be used with caution in individuals who are taking other drugs that may increase the risk of cancer, infection, or other serious conditions.



34575-50957

B/16A02B



QUALITY CHOICE



ENTERIC COATED

LO-DOSE ASPIRIN

ANALGESIC • PAIN RELIEVER • EASY TO SWALLOW

120 TABLETS 81 mg EACH

Complete
enteric coating
to allow
easy
swallowing



QUALITY CHOICE



ENTERIC COATED

LO-DOSE ASPIRIN

ANALGESIC • PAIN RELIEVER • EASY TO SWALLOW

120 TABLETS 81 mg EACH

Complete
enteric coating
to allow
easy
swallowing



ENTERIC COAT

LO-DO ASPIRIN

ANALGESIC PAIN RELIEVER • EASY TO

120 TABLETS 81 mg EACH



120 TABLETS 81 mg EACH
ANALGESIC • PAIN RELIEVER • EASY TO SWALLOW
ENTERIC COATED
LO-DOSE ASPIRIN

ACIDIC, WARNING: If you consume 3 or more
aspirin tablets every day, ask your doctor whether
you should use aspirin for pain relief.
Aspirin may irritate the stomach lining.
If you are taking a prescription drug for
heart or blood pressure, aspirin may
interfere with the action of these drugs.
Aspirin may also interfere with the action of
other drugs, such as blood thinners.
Aspirin may also interfere with the action of
other drugs, such as blood thinners.
Aspirin may also interfere with the action of
other drugs, such as blood thinners.

Aspirin may also interfere with the action of
other drugs, such as blood thinners.
Aspirin may also interfere with the action of
other drugs, such as blood thinners.
Aspirin may also interfere with the action of
other drugs, such as blood thinners.
Aspirin may also interfere with the action of
other drugs, such as blood thinners.

Aspirin may also interfere with the action of
other drugs, such as blood thinners.
Aspirin may also interfere with the action of
other drugs, such as blood thinners.
Aspirin may also interfere with the action of
other drugs, such as blood thinners.
Aspirin may also interfere with the action of
other drugs, such as blood thinners.



NDC 62770-227-12

FOR ARTHRITIS PAIN

ENTERIC COATED ASPIRIN

SAFETY COATED

SAFER FOR YOUR STOMACH THAN FLAVOIR OR BUTTERED ASPIRIN ANALGESIC

100 TABLETS-325 mg each



01021308



NDC 62770-227-12

FOR ARTHRITIS PAIN

ENTERIC COATED ASPIRIN

SAFETY COATED

SAFER FOR YOUR STOMACH THAN FLAVOIR OR BUTTERED ASPIRIN ANALGESIC

100 TABLETS-325 mg each



FOR ARTHRITIS PAIN

ENTERIC COATED ASPIRIN

100 TABLETS-325 mg each

Safer for your stomach than flavo or buttered aspirin.
INDICATIONS: For the temporary relief of minor aches and pains of arthritis and rheumatism. Enteric coated aspirin tablets are easier to swallow and are specially coated to help prevent the gastric upset often caused by plain uncoated aspirin.

DIRECTIONS: Adults and children over 12 years of age: 1 to 2 tablets every 4 hours, while symptoms persist, not to exceed 12 tablets in 24 hours, unless directed by a physician. Drink a full glass of water with each dose. Children under 12 years of age: Consult a physician.

WARNINGS: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye Syndrome, a rare but serious illness reported to be associated with aspirin. Do not take this product for pain for more than 10 days or for fever for more than 3 days unless directed

by a doctor. If pain or fever persists or gets worse, if new symptoms occur, or if redness or swelling is present, consult a doctor because these could be signs of a serious condition. Do not take this product if you are allergic to aspirin, have asthma, or if you have stomach problems that persist or recur, or if you have ulcers or bleeding problems unless directed by a doctor. If you get a hole or a hole of tearing occurs, consult a doctor before taking any more of this product. Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a poison control center immediately. As with any medicine, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. **IT IS ESPECIALLY IMPORTANT NOT TO USE ASPIRIN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS SPECIFICALLY DIRECTED TO DO SO BY A**

DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY.

ALCOHOL WARNING: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers. Aspirin may cause stomach bleeding.

DRUG INTERACTION PRECAUTIONS: Do not take this product if you are taking a prescription drug for anticoagulation (thinning of the blood), diabetes, gout or asthma unless directed by a physician.

Each tablet contains 325 mg Aspirin (81 mg USP) plus other inactive ingredients (may differ from brand): Docus Iron Oxide, Gelatin, Colloidal Silicon Dioxide, D&G Yellow #10 Aluminum Lake, FD&C Yellow #6 Aluminum Lake, Hydroxyethyl Methacrylate, Polyethylene Glycol, Polyvinyl Alcohol, Povidone, Polyethylene Glycol, Stearic Acid, Sodium Alginate, Sodium Bicarbonate,

Sorbitol, Sorbic Acid, Talc, Triacetin Dioxide, Triethyl Citrate.

Store in a cool, dry place at controlled room temperature 15°-30°C (59°-86°F). Use by expiration date on package. See end flap for expiration date and lot no.

TAMPER EVIDENT: DO NOT USE IF UNPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

* This product is not manufactured or distributed by Swearingin Research Corporation, owner of the registered trademark Control-T Tablets.

60834 Rev. 8/88
DISTRIBUTED BY:
STOP & SHOP
SUPERMARKET COMPANY
P.O. BOX 1042
BOSTON, MA 02101
MADE IN U.S.A.



This product can harm babies if swallowed. Keep out of reach of children.

The Medicine Shoppe

NDC 49114-117-72

Extra Strength Pain Reliever

Enteric Coated Aspirin

60 Tablets, 500 mg Each Safety Sealed

The Medicine Shoppe

NDC 49114-117-72

Extra Strength Pain Reliever

Enteric Coated Aspirin

60 Tablets, 500 mg Each Safety Sealed

The Medicine Shoppe

Extra Strength

Enteric Coated Aspirin

60 Tablets, 500 mg Each

B12212



Some use your stomach to relieve pain or to relieve aspirin.

INDICATIONS: For the temporary relief of minor aches and pains of arthritis and rheumatism. Enteric coated aspirin tablets are best to swallow and are specially coated to help prevent the gastric upset often caused by oral uncoated aspirin.

DIRECTIONS: ADULTS: 2 tablets every 8 hours, as necessary, with water or fruit juice. Do not exceed 8 tablets in 24 hours unless directed by a physician. CHILDREN under 12 years of age: As recommended by a physician.

WARNINGS: Children and teenagers should not use this medicine for chicken pox or flu symptoms unless a doctor is consulted. Avoid Reye Syndrome, a rare but serious illness reported to be associated with aspirin. Do not take this product for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor. If you or your pet have or had severe, if any symptoms such as, or if you notice or swelling is present, consult a doctor because these could be signs of a serious condition. Do not take this product if you are allergic to aspirin, have asthma, or if you have stomach problems that prevent or resist, or if you have ulcers or bleeding problems unless directed by a doctor. If dizziness or a loss of hearing occurs, a doctor helps taking any more of this product. Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a poison control center immediately. As with any medicine, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. **IT IS ESPECIALLY IMPORTANT NOT TO USE ASPIRIN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS SPECIFICALLY DIRECTED TO DO SO BY A DOCTOR.**

BECAUSE IT MAY CAUSE PROBLEMS IN THE UTERUS OR COMPLICATIONS DURING DELIVERY.

ACIDIC WARNING: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding.

DRUG INTERACTION PRECAUTION: Do not take this product if you are taking a medication for anticoagulation (thinning of blood), diabetes, joint or arthritis unless directed by a physician.

Each white coated tablet contains Aspirin 500 mg. May also contain (may differ from brand): Cellulose, Croscarmellose Sodium, Ethacrynic Acid, Polyvinyl Alcohol, Povidone, Talc, Polyethylene Glycol, Titanium Dioxide, Sodium Bicarbonate, Triethyl Citrate, Stearic Acid, Spoken Alginate, Colloidal Silicon Dioxide, Black Iron Oxide, Polyethylene Glycol, Hydroxypropyl Methylcellulose, Polysorbate, PO40 Yellow #6 Aluminum Lake, D & C Yellow #10 Aluminum Lake, Zinc Oxide.

Store in a cool, dry place at controlled room temperature 15° to 30°C (59° to 86°F). Use by expiration date on packaging. See use flap for expiration date and lot no.

TRAMPER EVIDENCE: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by SmithKline Beecham Consumer Brands, owner of the registered trademark Ecotrin® Aspirin Strength Tablets.

Made in USA 50844 1993 1298

Manufactured by
The Medicine Shoppe International, Inc.
St. Louis, MO 63137

SMITHKLINE BEECHAM
A GUARANTEE
OF THE HIGHEST QUALITY AND
STRENGTH OF OUR PRODUCTS
ON THE RECORD OF POSITIVE PROOF

018 30 1110 412 1 790 8201

CAP
EXTRA STRENGTH
PAIN RELIEVER
NON-ASPIRIN ANALGESIC
acetaminophen



DO NOT USE IF CAP IS BROKEN OR MISSING


CAP
EXTRA STRENGTH
PAIN RELIEVER
NON-ASPIRIN ANALGESIC
acetaminophen



DO NOT USE IF CAP IS BROKEN OR MISSING

CAP
EXTRA STRENGTH
PAIN RELIEVER
NON-ASPIRIN ANALGESIC
acetaminophen

DO NOT USE IF CAP IS BROKEN OR MISSING



Next
5

RETURN CARTON FOR COMPLETE PRODUCT INFORMATION

As with all other Extra Strength products, CAP is designed to relieve pain and reduce fever. It is not intended to cure any disease. CAP is a non-aspirin analgesic and antipyretic. It is not intended to be used for the relief of pain or fever in children under the age of 12 years. CAP is not intended to be used for the relief of pain or fever in patients with a history of liver disease or in patients with a history of alcohol abuse. CAP is not intended to be used for the relief of pain or fever in patients with a history of kidney disease or in patients with a history of heart disease. CAP is not intended to be used for the relief of pain or fever in patients with a history of diabetes or in patients with a history of high blood pressure. CAP is not intended to be used for the relief of pain or fever in patients with a history of asthma or in patients with a history of epilepsy. CAP is not intended to be used for the relief of pain or fever in patients with a history of glaucoma or in patients with a history of narrow-angle glaucoma. CAP is not intended to be used for the relief of pain or fever in patients with a history of bleeding disorders or in patients with a history of blood clotting disorders. CAP is not intended to be used for the relief of pain or fever in patients with a history of liver failure or in patients with a history of kidney failure. CAP is not intended to be used for the relief of pain or fever in patients with a history of heart failure or in patients with a history of high cholesterol. CAP is not intended to be used for the relief of pain or fever in patients with a history of diabetes or in patients with a history of high blood pressure. CAP is not intended to be used for the relief of pain or fever in patients with a history of asthma or in patients with a history of epilepsy. CAP is not intended to be used for the relief of pain or fever in patients with a history of glaucoma or in patients with a history of narrow-angle glaucoma. CAP is not intended to be used for the relief of pain or fever in patients with a history of bleeding disorders or in patients with a history of blood clotting disorders. CAP is not intended to be used for the relief of pain or fever in patients with a history of liver failure or in patients with a history of kidney failure. CAP is not intended to be used for the relief of pain or fever in patients with a history of heart failure or in patients with a history of high cholesterol.

Drug Facts for Extra Strength

Read the Drug Facts for Extra Strength carefully before you take it. Keep this information with the product. This information does not take the place of reading the label on the product. This information does not describe every possible use or warning. CAP is not intended to be used for the relief of pain or fever in children under the age of 12 years. CAP is not intended to be used for the relief of pain or fever in patients with a history of liver disease or in patients with a history of alcohol abuse. CAP is not intended to be used for the relief of pain or fever in patients with a history of kidney disease or in patients with a history of heart disease. CAP is not intended to be used for the relief of pain or fever in patients with a history of diabetes or in patients with a history of high blood pressure. CAP is not intended to be used for the relief of pain or fever in patients with a history of asthma or in patients with a history of epilepsy. CAP is not intended to be used for the relief of pain or fever in patients with a history of glaucoma or in patients with a history of narrow-angle glaucoma. CAP is not intended to be used for the relief of pain or fever in patients with a history of bleeding disorders or in patients with a history of blood clotting disorders. CAP is not intended to be used for the relief of pain or fever in patients with a history of liver failure or in patients with a history of kidney failure. CAP is not intended to be used for the relief of pain or fever in patients with a history of heart failure or in patients with a history of high cholesterol.

Directions

Adults and children 12 years of age and older: Take 1 tablet every 6 hours as needed. Do not use more than 4 tablets in 24 hours or 8 tablets in a day.

Children 12 years of age and older: Do not use. CAP is not intended to be used for the relief of pain or fever in children under the age of 12 years.

Other information

Keep this product out of the reach of children. CAP is not intended to be used for the relief of pain or fever in children under the age of 12 years. CAP is not intended to be used for the relief of pain or fever in patients with a history of liver disease or in patients with a history of alcohol abuse. CAP is not intended to be used for the relief of pain or fever in patients with a history of kidney disease or in patients with a history of heart disease. CAP is not intended to be used for the relief of pain or fever in patients with a history of diabetes or in patients with a history of high blood pressure. CAP is not intended to be used for the relief of pain or fever in patients with a history of asthma or in patients with a history of epilepsy. CAP is not intended to be used for the relief of pain or fever in patients with a history of glaucoma or in patients with a history of narrow-angle glaucoma. CAP is not intended to be used for the relief of pain or fever in patients with a history of bleeding disorders or in patients with a history of blood clotting disorders. CAP is not intended to be used for the relief of pain or fever in patients with a history of liver failure or in patients with a history of kidney failure. CAP is not intended to be used for the relief of pain or fever in patients with a history of heart failure or in patients with a history of high cholesterol.

Manufactured by: Watson Pharmaceuticals, Inc., Kalamazoo, MI 49001

Approved by: Consumer Safety Products, Inc., Tampa, FL 33603



Goldline

NDC 0182-1410-01 TAMPER-EVIDENT

GENAPAP™

acetaminophen (APAP) tablets.....ASPIRIN-free
PAIN RELIEVER/FEVER REDUCER 100 TABLETS
-325 mg EACH

Compare to the active ingredient in Regular Strength TYLENOL® Tablets

Goldline

NDC D182-141C-01 TAMPER-EVIDENT

GENAPAP™

acetaminophen (APAP) tablets.....ASPIRIN-free
PAIN RELIEVER/FEVER REDUCER 100 TABLETS
-325 mg EACH

Compare to the active ingredient in Regular Strength TYLENOL® Tablets

TAMPER-EVIDENT

Goldline

GENAPAP™

acetaminophen (APAP) tablets.....
ASPIRIN-free
PAIN RELIEVER/FEVER REDUCER
100 TABLETS - 325 mg EACH

Tamper-Evident package
See warning



Drug Facts (continued)	<p>Directions</p> <p>Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.</p> <p>Take 1 tablet every 4 to 6 hours as needed. Do not take more than 5 tablets in 24 hours.</p> <p>Do not drink alcohol. Do not use this with regular strength Tylenol in children under 6 years of age. This and other facts from the instructions list some important information you should know for your safety and health.</p> <p>Other Warnings: Do not use this with other products that contain acetaminophen. Do not use this with other products that contain aspirin. Do not use this with other products that contain salicylates. Do not use this with other products that contain NSAIDs. Do not use this with other products that contain opioids. Do not use this with other products that contain benzodiazepines. Do not use this with other products that contain sedatives. Do not use this with other products that contain anticholinergics. Do not use this with other products that contain MAO inhibitors. Do not use this with other products that contain SSRIs. Do not use this with other products that contain tricyclic antidepressants. Do not use this with other products that contain antipsychotics. Do not use this with other products that contain antiepileptics. Do not use this with other products that contain immunosuppressants. Do not use this with other products that contain antineoplastic agents. Do not use this with other products that contain antidiabetic agents. Do not use this with other products that contain antihypertensive agents. Do not use this with other products that contain anticholinergic agents. Do not use this with other products that contain antihistamines. Do not use this with other products that contain corticosteroids. Do not use this with other products that contain diuretics. Do not use this with other products that contain electrolyte supplements. Do not use this with other products that contain laxatives. Do not use this with other products that contain muscle relaxants. Do not use this with other products that contain antiarrhythmics. Do not use this with other products that contain antiplatelet agents. Do not use this with other products that contain anticoagulants. Do not use this with other products that contain anti-infective agents. Do not use this with other products that contain anti-parasitic agents. Do not use this with other products that contain anti-fungal agents. Do not use this with other products that contain anti-viral agents. Do not use this with other products that contain anti-cancer agents. Do not use this with other products that contain anti-aging agents. Do not use this with other products that contain anti-obesity agents. Do not use this with other products that contain anti-asthma agents. Do not use this with other products that contain anti-allergy agents. Do not use this with other products that contain anti-epileptic agents. Do not use this with other products that contain anti-seizure agents. Do not use this with other products that contain anti-spasmodic agents. Do not use this with other products that contain anti-convulsant agents. Do not use this with other products that contain anti-epileptic agents. Do not use this with other products that contain anti-seizure agents. Do not use this with other products that contain anti-spasmodic agents. Do not use this with other products that contain anti-convulsant agents.</p>
-------------------------------	---

This product has been tested and found to be free of opiates, barbiturates, and other addictive substances. Your doctor should be notified if you are taking any other drugs, especially if you are taking any of the following: alcohol, aspirin, salicylates, NSAIDs, opioids, benzodiazepines, sedatives, anticholinergics, MAO inhibitors, SSRIs, tricyclic antidepressants, antipsychotics, antiepileptics, immunosuppressants, antineoplastic agents, antidiabetic agents, antihypertensive agents, anticholinergic agents, antihistamines, corticosteroids, diuretics, electrolyte supplements, laxatives, muscle relaxants, antiarrhythmics, antiplatelet agents, anticoagulants, anti-infective agents, anti-parasitic agents, anti-fungal agents, anti-viral agents, anti-cancer agents, anti-aging agents, anti-obesity agents, anti-asthma agents, anti-allergy agents, anti-epileptic agents, anti-seizure agents, anti-spasmodic agents, anti-convulsant agents.

©2007 W. L. Bausch & Lomb, Inc. All rights reserved.
Distributed by: W. L. Bausch & Lomb, Inc., 1000 Lakeside Drive, Lake Bluff, IL 60044
Distributed by: W. L. Bausch & Lomb, Inc., 1000 Lakeside Drive, Lake Bluff, IL 60044

RETAIN CARD FOR COMPLETE PRODUCT INFORMATION

Drug Facts	Purpose Pain Reliever Fever Reducer
Active Ingredient (for each tablet)	Acetaminophen 325 mg
Uses: For the temporary relief of minor aches and pains associated with the flu, headache, muscular aches, or common cold. It is also used to reduce fever.	
Warnings: Do not exceed recommended dose. Avoid drinking alcohol or using alcohol. Do not use this with other products that contain acetaminophen. Do not use this with other products that contain aspirin. Do not use this with other products that contain salicylates. Do not use this with other products that contain NSAIDs. Do not use this with other products that contain opioids. Do not use this with other products that contain benzodiazepines. Do not use this with other products that contain sedatives. Do not use this with other products that contain anticholinergics. Do not use this with other products that contain MAO inhibitors. Do not use this with other products that contain SSRIs. Do not use this with other products that contain tricyclic antidepressants. Do not use this with other products that contain antipsychotics. Do not use this with other products that contain antiepileptics. Do not use this with other products that contain immunosuppressants. Do not use this with other products that contain antineoplastic agents. Do not use this with other products that contain antidiabetic agents. Do not use this with other products that contain antihypertensive agents. Do not use this with other products that contain anticholinergic agents. Do not use this with other products that contain antihistamines. Do not use this with other products that contain corticosteroids. Do not use this with other products that contain diuretics. Do not use this with other products that contain electrolyte supplements. Do not use this with other products that contain laxatives. Do not use this with other products that contain muscle relaxants. Do not use this with other products that contain antiarrhythmics. Do not use this with other products that contain antiplatelet agents. Do not use this with other products that contain anticoagulants. Do not use this with other products that contain anti-infective agents. Do not use this with other products that contain anti-parasitic agents. Do not use this with other products that contain anti-fungal agents. Do not use this with other products that contain anti-viral agents. Do not use this with other products that contain anti-cancer agents. Do not use this with other products that contain anti-aging agents. Do not use this with other products that contain anti-obesity agents. Do not use this with other products that contain anti-asthma agents. Do not use this with other products that contain anti-allergy agents. Do not use this with other products that contain anti-epileptic agents. Do not use this with other products that contain anti-seizure agents. Do not use this with other products that contain anti-spasmodic agents. Do not use this with other products that contain anti-convulsant agents.	

