

# Pharmaceutical Formulations

012 3059

13  
0000



# Allergy/Sinus Headache Caplets



Antihistamine  
Nasal Decongestant  
Pain Reliever

NDC # 10730-201-21

# Allergy/Sinus Headache Caplets

For the Temporary Relief of Nasal Congestion



Compare to active ingredients of  
Benadryl® Allergy Sinus Headache

24 CAPLETS



Allergy/Sinus  
Headache Caplets

EACH DOSE IS INDIVIDUALLY SEALED IN PLASTIC OVER A FOLDED BACKING.  
DO NOT USE IF THESE ARE FOUND OR PRESENT.

228117  
00014  
00000001-AM



0 72000 17560 6



100%  
Compare to active ingredients of  
Benadryl® Allergy Sinus Headache

Made in the USA  
100%  
Compare to active ingredients of  
Benadryl® Allergy Sinus Headache

DO NOT EXPOSE TO EXCESSIVE HEAT.  
NEVER IN A HOT PLACE.  
KEEP OUT OF REACH OF CHILDREN.  
SEEK MEDICAL ATTENTION IN  
CASE OF ACCIDENTAL OVER-  
DOSE. KEEP THIS AND ALL OTHERS OUT OF THE REACH  
OF CHILDREN. IN CASE OF ACCIDENTAL OVER-  
DOSE, CONTACT A POISON CONTROL CENTER OR  
YOUR PHYSICIAN IMMEDIATELY. IF YOU ARE  
PREGNANT OR BREASTFEEDING, CONSULT YOUR  
PHYSICIAN BEFORE USING THIS PRODUCT.

Indications: If you consume 2 or more  
tablets daily every day, ask your doctor whether  
you should have an examination or other tests done.  
Warnings: Do not use this product for more than  
10 days. If symptoms do not improve or are  
accompanied by fever, rash, or other signs of  
infection, stop use immediately and consult a doctor.  
Do not use if you have a history of asthma, or if you  
are taking any other medicine, or if you are taking  
any other medicine that may interact with this  
product. Do not use if you are taking any other  
medicine that may interact with this product.  
Do not use if you are taking any other  
medicine that may interact with this product.  
Do not use if you are taking any other  
medicine that may interact with this product.

Each Caplet Contains Three Active Ingredients:	First-Generation (Antihistamine) 10 mg (2 mg)
	Second-Generation (Antihistamine) 10 mg (2 mg)
	Third-Generation (Antihistamine) 10 mg (2 mg)
	Fourth-Generation (Antihistamine) 10 mg (2 mg)
	Fifth-Generation (Antihistamine) 10 mg (2 mg)
	Sixth-Generation (Antihistamine) 10 mg (2 mg)
	Seventh-Generation (Antihistamine) 10 mg (2 mg)
	Eighth-Generation (Antihistamine) 10 mg (2 mg)
	Ninth-Generation (Antihistamine) 10 mg (2 mg)
	Tenth-Generation (Antihistamine) 10 mg (2 mg)
	Eleventh-Generation (Antihistamine) 10 mg (2 mg)
	Twelfth-Generation (Antihistamine) 10 mg (2 mg)
	Thirteenth-Generation (Antihistamine) 10 mg (2 mg)
	Fourteenth-Generation (Antihistamine) 10 mg (2 mg)
	Fifteenth-Generation (Antihistamine) 10 mg (2 mg)
	Sixteenth-Generation (Antihistamine) 10 mg (2 mg)
	Seventeenth-Generation (Antihistamine) 10 mg (2 mg)
	Eighteenth-Generation (Antihistamine) 10 mg (2 mg)
	Nineteenth-Generation (Antihistamine) 10 mg (2 mg)
	Twentieth-Generation (Antihistamine) 10 mg (2 mg)
	Twenty-first-Generation (Antihistamine) 10 mg (2 mg)
	Twenty-second-Generation (Antihistamine) 10 mg (2 mg)
	Twenty-third-Generation (Antihistamine) 10 mg (2 mg)
	Twenty-fourth-Generation (Antihistamine) 10 mg (2 mg)
	Twenty-fifth-Generation (Antihistamine) 10 mg (2 mg)
	Twenty-sixth-Generation (Antihistamine) 10 mg (2 mg)
	Twenty-seventh-Generation (Antihistamine) 10 mg (2 mg)
	Twenty-eighth-Generation (Antihistamine) 10 mg (2 mg)
	Twenty-ninth-Generation (Antihistamine) 10 mg (2 mg)
	Thirtieth-Generation (Antihistamine) 10 mg (2 mg)



0100001248



0100001248

9  
AUG  
1998

LOT:  
EXP:

# MAXIMUM STRENGTH NON-ASPIRIN FLU MEDICINE



**90** 100% SATISFACTION GUARANTEED  
*Guarantee*

**DOLLAR  
GENERAL**  
**\$1.50**  
DEPT. D  
EVERY DAY

MOE 55110 410 10

MAXIMUM STRENGTH  
NON-ASPIRIN

# FLU MEDICINE

Compare to the  
active ingredients  
of Tylenol Flu

FOR THE NON-DROWSY TEMPORARY REMEDIES

- Fever
- Headaches / Body Aches
- Coughing
- Nasal Congestion
- Sore Throat

**10 GELATIN CAPLETS**

254150  
SP10  
EXAM16 OF

# MAXIMUM STRENGTH NON-ASPIRIN FLU MEDICINE



08604 19107









1 UNIT



Non-Aspirin Maximum Strength  
**Pain Reliever  
Sinus/Allergy**



LOT:  
EXP:



Non-Aspirin Maximum Strength

# Pain Reliever Sinus/Allergy



EACH BOX IS INDIVIDUALLY SEALED  
IN PLASTIC OVER A FOIL BACKING.  
DO NOT USE IF THESE ARE TORN OR BROKEN.



Antihistamine • Nasal Decongestant  
For the Temporary Relief of Pain, Fever,  
Fchy, Watery Eyes, Nasal Congestion

Compare To The Active Ingredients  
Tylenol® Allergy Sinus\*

24 Caplets

Non-Aspirin Maximum Strength  
**Pain Reliever  
Sinus/Allergy**



2007  
MAY 31  
BATCH OF 44



1-800-432-6111

**Substitution Guaranteed**  
MAY BE SUBSTITUTED WITH AN EQUIVALENT BRAND OF THE SAME CLASS AND STRENGTH.  
DO NOT USE IF THESE ARE TORN OR BROKEN.  
EACH BOX IS INDIVIDUALLY SEALED IN PLASTIC OVER A FOIL BACKING.  
DO NOT USE IF THESE ARE TORN OR BROKEN.  
EACH BOX IS INDIVIDUALLY SEALED IN PLASTIC OVER A FOIL BACKING.  
DO NOT USE IF THESE ARE TORN OR BROKEN.



**ACTIVE INGREDIENTS:** Oxycodone Hydrochloride 5 mg, Acetaminophen 325 mg  
**INACTIVE INGREDIENTS:** Cellulose, Dextrose, Hydroxypropyl Cellulose, Magnesium Stearate, Polyethylene Glycol, Talc, Triacetin, Yellow Iron Oxide  
**HOW TO USE:** Take 2 caplets 4 times a day with food or milk. Do not exceed 12 caplets in 24 hours.  
**CAUTION:** Do not drink alcohol while taking this medicine. Do not operate machinery or drive a car until you know how this medicine affects you.  
**WARNING:** This medicine may cause drowsiness. Do not take with other medicines that cause drowsiness.  
**PRECAUTIONS:** Tell your doctor if you are taking any other medicines, especially other pain relievers, sedatives, tranquilizers, or sleeping pills.  
**CONTRAINDICATIONS:** Do not take if you are allergic to oxycodone, acetaminophen, or any of the ingredients.  
**ADVERSE REACTIONS:** Drowsiness, constipation, dry mouth, nausea, vomiting, dizziness, headache, blurred vision, difficulty breathing, and changes in heart rate.  
**DRUG INTERACTIONS:** This medicine may interact with other medicines, including alcohol, sedatives, tranquilizers, and sleeping pills.  
**PREGNANCY AND LACTATION:** Tell your doctor if you are pregnant, planning to get pregnant, or are breastfeeding.  
**USE IN CHILDREN:** Do not use in children under 18 years of age.  
**USE IN ELDERLY:** Use with caution in the elderly.  
**OVERDOSE:** If you take too much, you may experience symptoms such as drowsiness, confusion, and difficulty breathing. Call your doctor or poison control center immediately.  
**KEEP OUT OF REACH OF CHILDREN.**











